

Setup for the Josephson effect

In this section we describe in greater detail the setup we considered in the linear response calculation, relaxing some of the conditions assumed in the article. Although we use the notation and terminology of Figure 1a of the main article the description is equivalent for example with the system of Figure 1b of the main article.

In order to accommodate a superfluid with balanced pairing, we require a balanced number of particles within the superfluid states *i.e.* $N_1 = N_2$ and $N_3 = N_4$. We further assume that for any tunneling in the system the relation $\delta N_i/N_i \ll 1$ holds, so that we do not have to take into account the internal dynamics of the superfluids. In particular, this assumption implies that the superfluid states are not destroyed because of the transitions. The tunneling link between the hyperfine states is provided by two radio frequency (RF) fields $\mathbf{E}_{13}(\mathbf{r}, t)$ and $\mathbf{E}_{24}(\mathbf{r}, t)$. The field $\mathbf{E}_{ij}(\mathbf{r}, t)$ drives transitions between the states $|i\rangle$ and $|j\rangle$. For simplicity we take these fields as a single mode with frequency ν_{ij} and wave vector $\mathbf{k}_{ij}$. The strength of the coupling between the electric field $\mathbf{E}_{ij}(\mathbf{r}, t)$ and the hyperfine transition between states $|i\rangle$ and $|j\rangle$ is given by the Rabi frequency

$$\Omega_{ij}(\mathbf{r}) = \mathbf{d}_{ij} \cdot \mathbf{E}_{ij}(\mathbf{r}), \quad (1)$$

where $\mathbf{d}_{ij}$ is the electric dipole moment of the transition. Throughout the treatment we assume that the RF fields are near resonance and that the intensity of the radiation is low. With these assumptions we may use the rotating wave approximation [1]. As a result, it is possible to transform the time dependence of the field in the rotating frame into a shift in the transition frequency δ_{ij} , called detuning, defined as

$$\delta_{ij} = \nu_{ij} - \omega_{ij}, \quad (2)$$

where ω_{ij} is the frequency of the transition.

Within the same superfluid state the hyperfine states will naturally have the same chemical potential, therefore $\mu_1 = \mu_2$ and $\mu_3 = \mu_4$. In the article we further assumed $\mu_1 = \mu_3$, but here we remove this constraint. Including this difference in the chemical potential between the two superfluids into the RF detuning, we get the modified detuning of the hyperfine transition

$$\tilde{\delta}_{ij} = \delta_{ij} + \mu_j - \mu_i. \quad (3)$$

This quantity describes the energy gain or loss associated with a particular hyperfine transition without many-body effects.




Figure 1: The energy levels of the system. In the unperturbed system the chemical potentials are equal within the BCS state: $\mu_1 = \mu_2$ and $\mu_3 = \mu_4$. The detunings δ_{13} and δ_{24} of the RF fields can create an effective asymmetry in the energies.

The energy levels of the system in question are shown in Figure 1. Here we make use of the assumption that the applied RF field does not alter the internal dynamics of the superfluids so that we may split the detuning δ_{ij} between the states $|i\rangle$ and $|j\rangle$ freely. We choose a symmetric splitting, although we will eventually see that this choice does not affect the results.

Linear response description

The system is described by the Hamiltonian

$$H = H_0 + H_{RF}. \quad (4)$$

Here we have defined H_0 as

$$H_0 = \int d\mathbf{r} \sum_i \psi_i^\dagger(\mathbf{r}) \left(-\frac{\nabla^2}{2m} - \mu_i \right) \psi_i(\mathbf{r}) + \frac{1}{2} \int d\mathbf{r} \sum_{i \neq j} U_{ij} \psi_i^\dagger(\mathbf{r}) \psi_j^\dagger(\mathbf{r}) \psi_j(\mathbf{r}) \psi_i(\mathbf{r}). \quad (5)$$

The external perturbation is given by

$$H_{RF} = H_c + H_d, \quad (6)$$

where we have

$$H_c = \int d\mathbf{r} \sum_{i,j} \Omega_{ij} \psi_i^\dagger(\mathbf{r}) \psi_j(\mathbf{r}) \quad (7)$$

$$H_d = \int d\mathbf{r} \sum_i W_i \psi_i^\dagger(\mathbf{r}) \psi_i(\mathbf{r}), \quad (8)$$

with the definitions $W_1 = \mu_1 + \delta_{13}/2$, $W_3 = \mu_3 - \delta_{13}/2$ and similarly for W_2 and W_4 . The inclusion of the chemical potential in these expressions is just a mathematical trick to aid the forthcoming calculation. For RF fields it is a very good approximation to take the Rabi frequency Ω as independent of position.

The operator for current density to one hyperfine state is given by the time derivative of the particle density operator, which is obtained using the Heisenberg equation of motion

$$\frac{d}{dt} N_i(\mathbf{r}_1) = i [H, N_i(\mathbf{r}_1)]. \quad (9)$$

Only H_c fails to commute with the number operator. Hence the Heisenberg equation of motion for the number operator becomes

$$\begin{aligned} \frac{d}{dt} N_i(\mathbf{r}_1) &= i [H_c(t), N_i(\mathbf{r}_1)] \\ &= i \sum_j \int d\mathbf{r}_2 \left(\Omega_{ji} \psi_j^\dagger(\mathbf{r}_2) \psi_i(\mathbf{r}_1) - \Omega_{ij} \psi_i^\dagger(\mathbf{r}_1) \psi_j(\mathbf{r}_2) \right). \end{aligned} \quad (10)$$

The physical current density $I_i(\mathbf{r}_1)$ is the thermodynamic average of the current density operator defined as $I_i(\mathbf{r}_1 t) \equiv \langle \dot{N}_i(\mathbf{r}_1) \rangle$, which we then need to calculate. However, the exact calculation of this quantity is not feasible, since time evolution of the operators in the Heisenberg picture depends on the coupling H_c . Hence, we need to resort to an approximate solution with respect to the coupling strength. At this point it is most convenient to make a transformation to the interaction picture with respect to H_c . For a general operator $A(t)$ this transformation is given by

$$A(t) = S(t, t_0)^{-1} A(t_0) S(t, t_0), \quad (11)$$

in which t_0 gives the initial condition and

$$S(t, t_0) = T \exp \left(-i \int_{t_0}^t dt' H_c(t') \right). \quad (12)$$

Assuming that H_c is a weak perturbation, it is a good approximation to linearise this transformation with respect to H_c , in which case we have

$$S(t, t_0) = 1 - i \int_{t_0}^t dt' H_c(t'), \quad (13)$$

and consequently $A(t)$ is linearised to

$$A(t) = A(t_0) - i \int_{t_0}^t dt' [A(t'), H_c(t')]. \quad (14)$$

The thermodynamic expectation value of a generic operator $A(t)$ then follows the Kubo formula

$$\langle A(t) \rangle = \langle A(t_0) \rangle - i \int_{t_0}^t dt' \langle [A(t'), H_c(t')] \rangle. \quad (15)$$

Here t_0 is the reference time for the initial condition. Since we assume a system initially in equilibrium, there is no current at the time t_0 . Therefore, upon replacing A with the current operator $\hat{N}$, the first term in the expression (15) vanishes. In order to obtain the Kubo formula in the frequency domain, it is a common strategy to take the limit $t_0 \rightarrow -\infty$ for the initial condition.

Inserting equation (10) into equation (15) then yields the current density

$$\begin{aligned} I_i(\mathbf{r}_1 t) = & i \int_{-\infty}^{\infty} dt' \sum_{j,k,l} \int d\mathbf{r}_2 d\mathbf{r}_3 d\mathbf{r}_4 \left(\exp[i(W_l - W_i)t + i(W_j - W_k)t'] \right. \\ & \left. \times \Omega_{li} \Omega_{jk} L_{ijkl}(\mathbf{r}_1 t, \mathbf{r}_3 t, \mathbf{r}_4 t', \mathbf{r}_2 t') - h.c. \right), \end{aligned} \quad (16)$$

where the retarded linear response function $L_{ijkl}(\mathbf{r}_1 t, \mathbf{r}_2 t, \mathbf{r}_3 t', \mathbf{r}_4 t')$ is defined as

$$L_{ijkl}(\mathbf{r}_1 t, \mathbf{r}_2 t, \mathbf{r}_3 t', \mathbf{r}_4 t') = -i\theta(t - t') \left\langle \left[\psi_k^\dagger(\mathbf{r}_3 t) \psi_i(\mathbf{r}_1 t), \psi_l^\dagger(\mathbf{r}_4 t') \psi_j(\mathbf{r}_2 t') \right] \right\rangle. \quad (17)$$

In equation (16) we have used standard algebraic manipulations to separate out the effect of the quantities W_i from the remaining time evolution of the field operators. For the hyperfine species $|1\rangle$, equation (16) can be rewritten as

$$I_{13}(\mathbf{r}_1 t) = I_{13}^S(\mathbf{r}_1 t) + I_{13}^J(\mathbf{r}_1 t), \quad (18)$$

where the standard single particle (quasiparticle) current I_{13}^S is defined as

$$I_{13}^S(\mathbf{r}_1 t) = -2 \int d\mathbf{r}_2 \text{Im} \left[\Omega_{13}^* \Omega_{13} L_{1331}(\mathbf{r}_1 - \mathbf{r}_2, \tilde{\delta}_{13} + i\eta^+) \right], \quad (19)$$

and the Josephson current I_{13}^J as

$$\begin{aligned} I_{13}^J(\mathbf{r}_1 t) = & -2 \int d\mathbf{r}_2 \text{Im} \left[e^{-i(\tilde{\delta}_{13} + \tilde{\delta}_{24})t} \Omega_{13}^* \Omega_{24}^* \right. \\ & \left. \times L_{1234}(\mathbf{r}_1 - \mathbf{r}_2, -\tilde{\delta}_{24} + i\eta^+) \right]. \end{aligned} \quad (20)$$

In Eqs. (19, 20) we have identified the Fourier transformation in time for L . This was the reason for separating the quantities W_i from the time-evolution. In order to guarantee the convergence of the transformation, an infinitesimal (positive) convergence factor η^+ has been introduced. Thus, we see that the modified detunings defined already in equation (3) now appear in the expression for the current. Here we may identify the spatial integration as a Fourier transformation with respect to zero momentum. Furthermore, in a homogeneous system we may drop the variable $\mathbf{r}_1$ since the current is not a function of position. Therefore we obtain

$$\begin{aligned} I_{13}^S(t) = & -2 \text{Im} \left[\Omega_{13}^* \Omega_{13} L_{1331}(\mathbf{p} = 0, \tilde{\delta}_{13} + i\eta^+) \right], \\ I_{13}^J(t) = & -2 \text{Im} \left[e^{-i(\tilde{\delta}_{13} + \tilde{\delta}_{24})t} \Omega_{13}^* \Omega_{24}^* L_{1234}(\mathbf{p} = 0, -\tilde{\delta}_{24} + i\eta^+) \right]. \end{aligned} \quad (21)$$

We note here that the definitions of single-particle and Josephson currents agree with their definitions in the conventional case of two superconductors when $\tilde{\delta}_{13} = \tilde{\delta}_{24}$.

At this point in our calculation we have isolated the effect of the perturbation on the initial system in the linear response approximation. Hereafter we only need to work with linear response functions which depend only on the properties of the unperturbed system.

Kadanoff-Baym formalism

Our goal here is the calculation of the retarded linear response function L . To this aim we will resort to the Kadanoff-Baym formalism [2, 3] to which we refer also as the self-consistent method. The interest of this method for our purposes lies in the fact that it allows the inclusion of the cross-interaction effects in the transport properties, and that, on more general grounds, the linear response function L will obey the same conservation laws that are obeyed by the single-particle Green's function $\mathcal{G}$.

In the Kadanoff-Baym formalism, when the general case of a time and position dependent external perturbation is considered, L is obtained from $\mathcal{G}$ by functional differentiation with respect to the external field

$$L_{ijkl}(\mathbf{r}_i\tau, \mathbf{r}_j\tau, \mathbf{r}_k\tau', \mathbf{r}_l\tau') = \left[\frac{\delta \mathcal{G}_{ik}(\mathbf{r}_i\tau, \mathbf{r}_k\tau')}{\delta \Omega_{lj}(\mathbf{r}_l\tau', \mathbf{r}_j\tau)} \right]_{\Omega=0}. \quad (22)$$

Here we are working in complex time Matsubara formalism, which is a standard technique for dealing with finite temperature Green's functions. The real-time retarded linear response function can be obtained from L in the Matsubara formalism by analytical continuation.

For the sake of brevity, we truncate the variables such as $\mathbf{r}_i$ and the spin-index in the following two equations to just the index i . Furthermore, we denote the variables of integration with a bar over the variable. The essential starting point is the equation of motion for the single particle Green's function

$$\int \mathcal{G}_0^{-1}(1, \bar{1}) \mathcal{G}(\bar{1}, 1') = \delta(1 - 1') + \int \Omega(1, \bar{1}) \mathcal{G}(\bar{1}, 1') + \int \Sigma(1, \bar{1}) \mathcal{G}(\bar{1}, 1'), \quad (23)$$

Here $\mathcal{G}_0^{-1}$ is the inverse free propagator, which accounts for the kinetic part of the equation of motion, and the self-energy Σ contains all the interaction effects in the system. On the level of principle one should solve this equation for an arbitrary perturbation Ω to obtain $\mathcal{G}$ and then further calculate L . This is in practice unfeasible, and in the Kadanoff-Baym method one circumvents this problem by taking the variational derivative of the equation of motion at $\Omega = 0$. This leads to an implicit equation for L

$$\begin{aligned} L(12, 1'2') &= \int \mathcal{G}(1, \bar{3})_{\Omega=0} \mathcal{G}(\bar{4}, 1')_{\Omega=0} \left[\frac{\delta \Omega(\bar{3}, \bar{4})}{\delta \Omega(2', 2)} \right]_{\Omega=0} \\ &+ \int \mathcal{G}(1, \bar{3})_{\Omega=0} \mathcal{G}(\bar{4}, 1')_{\Omega=0} \left[\frac{\delta \Sigma(\bar{3}, \bar{4})}{\delta \mathcal{G}(\bar{5}, \bar{6})} \right]_{\Omega=0} L(\bar{5}2, \bar{6}2'), \end{aligned} \quad (24)$$

which is easier to solve. In fact, in a homogeneous system we may utilize Fourier transformations leading to a matrix equation for L in momentum space.

In the following calculation we consider the case where $U_{13} \neq 0$; there is no conceptual difference when also other cross interactions are considered different from zero. If the BCS approximation is considered for the unperturbed system, the explicit form of the linear response function becomes

$$\begin{aligned} L_{1331}(\mathbf{p}, \omega) &= \frac{\Pi_{\mathcal{G}}(\mathbf{p}, \omega)}{1 + U_{13}\Pi_{\mathcal{G}}(\mathbf{p}, \omega)}, \\ L_{1234}(\mathbf{p}, \omega) &= -\frac{\Pi_{\mathcal{F}}(\mathbf{p}, \omega)}{1 + U_{13}\Pi_{\mathcal{G}}(\mathbf{p}, \omega)} \end{aligned} \quad (25)$$

for the current between states $|1\rangle$ and $|3\rangle$ and

$$\begin{aligned} L_{2442}(\mathbf{p}, \omega) &= \Pi_{\mathcal{G}}(\mathbf{p}, \omega) - \frac{U_{13}\Pi_{\mathcal{F}}(\mathbf{p}, \omega)^2}{1 + U_{13}\Pi'_{\mathcal{G}}(\mathbf{p}, \omega)}, \\ L_{2143}(\mathbf{p}, \omega) &= -\frac{\Pi_{\mathcal{F}}(\mathbf{p}, \omega)}{1 + U_{13}\Pi'_{\mathcal{G}}(\mathbf{p}, \omega)} \end{aligned} \quad (26)$$

for the current between states $|2\rangle$ and $|4\rangle$.

The terms Π in the previous equations are

$$\Pi_{\mathcal{G}}(\mathbf{p}, \omega) = \frac{1}{\beta V} \sum_{\mathbf{q}, \chi} \mathcal{G}_{11}(\mathbf{q}, \chi) \mathcal{G}_{33}(\mathbf{q} - \mathbf{p}, \chi - \omega), \quad (27)$$

$$\Pi'_{\mathcal{G}}(\mathbf{p}, \omega) = \frac{1}{\beta V} \sum_{\mathbf{q}, \chi} \mathcal{G}_{33}(\mathbf{q}, \chi) \mathcal{G}_{11}(\mathbf{q} - \mathbf{p}, \chi - \omega), \quad (28)$$

$$\Pi_{\mathcal{F}}(\mathbf{p}, \omega) = \frac{1}{\beta V} \sum_{\mathbf{q}, \chi} \mathcal{F}_{12}(\mathbf{q}, \chi) \mathcal{F}_{34}^*(\mathbf{q} - \mathbf{p}, \chi - \omega). \quad (29)$$

Here $\mathcal{F}$ is the so called anomalous Green's function, which expresses the pair correlation of the superfluid. In the main article we study a case where the two superfluids are identical, in which case we have $\Pi_{\mathcal{G}} = \Pi'_{\mathcal{G}}$.

The expression for the linear response function leads to the following results for the single-particle and Josephson currents

$$I_{13}^S(t) = -2|\Omega_{13}|^2 \times \text{Im} \left[\frac{\Pi_{\mathcal{G}}(\mathbf{p} = 0, \tilde{\delta}_{13} + i\eta^+)}{1 + U_{13}\Pi_{\mathcal{G}}(\mathbf{p} = 0, \tilde{\delta}_{13} + i\eta^+)} \right], \quad (30)$$

$$I_{24}^S(t) = -2|\Omega_{24}|^2 \times \text{Im} \left[\Pi_{\mathcal{G}}(\mathbf{p} = 0, \tilde{\delta}_{24} + i\eta^+) - \frac{U_{13}\Pi_{\mathcal{F}}(\mathbf{p} = 0, \tilde{\delta}_{24} + i\eta^+)^2}{1 + U_{13}\Pi'_{\mathcal{G}}(\mathbf{p} = 0, \tilde{\delta}_{24} + i\eta^+)} \right]. \quad (31)$$

$$I_{13}^J(t) = 2 \left| \Omega_{13}\Omega_{24} \frac{\Pi_{\mathcal{F}}(\mathbf{p} = 0, -\tilde{\delta}_{24} + i\eta^+)}{1 + U_{13}\Pi_{\mathcal{G}}(\mathbf{p} = 0, -\tilde{\delta}_{24} + i\eta^+)} \right| \times \sin[(\tilde{\delta}_{13} + \tilde{\delta}_{24})t + \varphi(\tilde{\delta}_{24})], \quad (32)$$

$$I_{24}^J(t) = 2 \left| \Omega_{13}\Omega_{24} \frac{\Pi_{\mathcal{F}}(\mathbf{p} = 0, -\tilde{\delta}_{13} + i\eta^+)}{1 + U_{13}\Pi'_{\mathcal{G}}(\mathbf{p} = 0, -\tilde{\delta}_{13} + i\eta^+)} \right| \times \sin[(\tilde{\delta}_{13} + \tilde{\delta}_{24})t + \varphi(\tilde{\delta}_{13})]. \quad (33)$$

The phase factors $\varphi(\tilde{\delta}_{13})$ and $\varphi(\tilde{\delta}_{24})$ which appear in equations (32) and (33) are the initial complex phase of the retarded linear response function L . In the article we have left out the minus sign from the detunings in the terms Π since in the case of identical superfluids, which we consider in the article, the critical current becomes a symmetric function of the detuning.

Identical superfluids

Here we discuss the results obtained in the case of identical superfluids. In this case the notation simplifies as $\tilde{\delta}_{13} = \delta_{13} + \mu_3 - \mu_1 = \delta_{13}$ and $\tilde{\delta}_{24} = \delta_{24} + \mu_4 - \mu_2 = \delta_{24}$ since the chemical potentials are equal. From the Kadanoff-Baym formalism, in the case where cross-interactions are zero, we have

$$\begin{aligned} I_{13}^S(\mathbf{r}_1 t) &= -2|\Omega_{13}|^2 \text{Im} [\Pi_{\mathcal{G}}(\mathbf{p} = 0, \delta_{13} + i\eta^+)], \\ I_{24}^S(\mathbf{r}_1 t) &= -2|\Omega_{24}|^2 \text{Im} [\Pi_{\mathcal{G}}(\mathbf{p} = 0, \delta_{24} + i\eta^+)], \end{aligned} \quad (34)$$

and

$$\begin{aligned} I_{13}^J(\mathbf{r}_1 t) &= 2 |\Omega_{13}\Omega_{24}\Pi_{\mathcal{F}}(\mathbf{p} = 0, -\delta_{24} + i\eta^+)| \sin((\delta_{13} + \delta_{24})t + \varphi_1(\delta_{24})), \\ I_{24}^J(\mathbf{r}_1 t) &= 2 |\Omega_{13}\Omega_{24}\Pi_{\mathcal{F}}(\mathbf{p} = 0, -\delta_{13} + i\eta^+)| \sin((\delta_{13} + \delta_{24})t + \varphi_2(\delta_{13})). \end{aligned} \quad (35)$$

The single-particle current vanishes when $|\delta_{ij}| < 2\Delta_{12}$ (Fig. 2). In this range only the Josephson current exists, as it is intuitively clear, since the single-particle current requires the breaking of Cooper pairs. At $|\delta_{ij}| = 2\Delta_{12}$ the detuning equals the minimum of the transition energy $E_{12}(\mathbf{q}) + E_{34}(\mathbf{q})$, where $E_{\alpha\beta}(\mathbf{q}) = \sqrt{(|q|^2/2m - \mu_{\alpha})^2 + |\Delta_{\alpha\beta}|^2}$.

This occurs at the point $\frac{q^2}{2m} = \mu$ *i.e.* at the Fermi level, where the BCS density of states is divergent, leading to a divergence in the dependence of the currents on the detuning. In the case of the critical Josephson current, this divergence is commonly known as the Riedel peak.

To illustrate the spin-asymmetry further, consider the results of Fig. 2. Notice that we plot only I_{13}^C since I_{24}^C is the same function, the difference being only that I_{13}^C depends on δ_{24} whereas I_{24}^C depends on δ_{13} . For instance




Figure 2: (left to right). The single-particle current I_{13}^S and the critical Josephson current I_{13}^C vs. δ_{13} and δ_{24} for identical BCS states. $U_{12} = U_{34} = -2.0E_F/n$. T is in the units of Fermi temperature T_F and in the temperature range of the figure, the gap is $\Delta \sim 0.4E_F$.




Figure 3: (left to right). The single-particle current I_{13}^S and the critical Josephson current I_{13}^C vs. $\tilde{\delta}_{13}$ and $\tilde{\delta}_{24}$ for unidentical BCS states. $U_{12} = -1.0E_F/n$, $U_{34} = -5.0E_F/n$, T is in the units of Fermi temperature T_F .

taking $\delta_{13}/E_F = 0.4$ and $\delta_{24}/E_F = 0.5$ at $T/T_F = 0.08$ then leads to $I_{13}^C(\delta_{24})/I_{24}^C(\delta_{13}) = 1.14$, in other words, a 14 % difference in the amplitude of the Josephson current for each spin component.

Finite temperature and cross interactions

It is instructive and of certain interest, from the experimental point of view, to discuss how the results presented in the article are affected by temperature and different interaction strengths. In order to properly observe finite temperature effects, we consider the slightly more complicated case of two different BCS states. Experimentally, the condensates 1-2 and 3-4 may have different interaction strengths. We choose $U_{12} = -1.0E_F/n$ and $U_{34} = -5.0E_F/n$. This leads to gaps $\Delta_{12} \sim 0.2E_F$ and $\Delta_{34} \sim 0.5E_F$, which corresponds roughly to systems that have been realised experimentally. Also the choice of temperature we use, $T = 0.06 \dots 0.1T_F$, is motivated by the lowest temperatures reached in the experiments. The results are plotted in Figs. 3 and 4. First of all we notice that all the features previously found still dominate the result, even though there are some clear changes.

We see that for finite T and $U_{12} \neq U_{34}$ there can be a finite single particle current even at small detunings. This current arises due to thermal excitations in the BCS states. Intuitively, this contribution is generated by Cooper pairs that have been already broken because of thermal energy. To be more precise, it is the fact that one of the BCS states has more thermal excitations for a given energy than the other that leads to the emergence of the thermal single particle current. We did indeed have thermal excitations in the results of the previous section as well, but their net contribution summed up to zero. The new peaks in the results (see Figures 3 and 4) are associated with the the contribution of thermal excitations diverging at a particular detuning. The explanation for these divergences is similar as for the divergences encountered in the previous section. Here the energy of the tunneling process is $E_{12}(\mathbf{q}) - E_{34}(\mathbf{q})$ (again with $E_{\alpha\beta}(\mathbf{q}) = \sqrt{(\frac{1}{2m}|\mathbf{q}|^2 - \mu_\alpha)^2 + |\Delta_{\alpha\beta}|^2}$) and the resonance occurs when the detuning coincides with the extremum of this energy. The density of states argument of the previous section holds then here as well. Moreover, the thermal single particle current vanishes above this maximum because at larger detunings there are no thermal transitions that would conserve both energy and momentum.




Figure 4: Single-particle current I_{13}^S , and critical Josephson current I_{13}^C vs. $\tilde{\delta}_{13}$ and $\tilde{\delta}_{24}$ for unidentical BCS states. $U_{12} = -1.0E_F/n$, $U_{34} = -5.0E_F/n$, T is in the units of Fermi temperature T_F .




Figure 5: Energy levels for the four-state model.

The perturbative calculation for the four-state system

The Hamiltonian

For the perturbative calculation of the four-state system of Figure 5, we consider the following Hamiltonian

$$H = H_0 + H_{RF}, \quad (36)$$

where H_0 contains the energies of the states and H_{RF} is the RF coupling. They have the form

$$H_0 = -U (n_1 n_2 + n_3 n_4),$$

$$H_{RF} = \Omega_{13} c_1^\dagger c_3 + h.c. + \Omega_{24} c_2^\dagger c_4 + h.c. + \frac{\delta_{13}}{2} (n_1 - n_3) + \frac{\delta_{24}}{2} (n_2 - n_4), \quad (37)$$

where U is the energy difference between the paired and unpaired states. We have kept the notation here as similar as possible to the initial description in the article. Obviously one could reformulate this four-state system in a more compact form.

Perturbative calculation

Now let us study our model system analytically using perturbation theory. The dynamics of the system can be described by four states: the paired states $|\phi_I\rangle \equiv |12\rangle$, $|\phi_{II}\rangle \equiv |34\rangle$ – corresponding to the left and right well respectively in the double well description – and the states $|14\rangle$, $|23\rangle$ (see Figure 5).

In order to find the lowest order contribution of the RF-couplings we have to employ the time-dependent perturbation theory to the second order in the RF-couplings. As an initial state we consider an arbitrary superposition of the paired states

$$|\phi_0\rangle = \alpha_0 |\phi_I\rangle + \beta_0 |\phi_{II}\rangle. \quad (38)$$

We point out already that in the $\alpha_0 = 0$ or $\beta_0 = 0$ case no Josephson current will appear. Calculating the occupation number N_1 of the hyperfine state in the state $|\phi(t)\rangle = \exp[-iHt]|\phi_0\rangle$ gives

$$\begin{aligned} \langle\phi(t)|N_1|\phi(t)\rangle &= |\alpha_0|^2 \langle\phi_I(t)|N_1|\phi_I(t)\rangle + |\beta_0|^2 \langle\phi_{II}(t)|N_1|\phi_{II}(t)\rangle \\ &+ \alpha_0\beta_0^* \langle\phi_{II}(t)|N_1|\phi_I(t)\rangle + \alpha_0^*\beta_0 \langle\phi_I(t)|N_1|\phi_{II}(t)\rangle. \end{aligned} \quad (39)$$

The perturbative expansion for $|\phi_I(t)\rangle$ and $|\phi_{II}(t)\rangle$ reads

$$\begin{aligned} |\phi_I(t)\rangle &= |12\rangle + \gamma_{12}^{(1)}(t)|14\rangle + \delta_{12}^{(1)}(t)|23\rangle + [\alpha_{14\,12}^{(2)}(t) + \alpha_{23\,12}^{(2)}(t)]|12\rangle + [\beta_{14\,12}^{(2)}(t) + \beta_{23\,12}^{(2)}(t)]|34\rangle \\ |\phi_{II}(t)\rangle &= |34\rangle + \gamma_{34}^{(1)}(t)|14\rangle + \delta_{34}^{(1)}(t)|23\rangle + [\alpha_{14\,34}^{(2)}(t) + \alpha_{23\,34}^{(2)}(t)]|12\rangle + [\beta_{14\,34}^{(2)}(t) + \beta_{23\,34}^{(2)}(t)]|34\rangle \end{aligned} \quad (40)$$

in equation (40) the superscript represents the perturbative order, the subscript the path corresponding to the process considered and the Greek letter its starting state (with the convention $\alpha \leftrightarrow |12\rangle$, $\beta \leftrightarrow |34\rangle$, $\gamma \leftrightarrow |14\rangle$, $\delta \leftrightarrow |23\rangle$). For instance $\beta_{14\,12}^{(2)}$ corresponds to the second-order process $|34\rangle \rightarrow |14\rangle \rightarrow |12\rangle$. The coefficients that we need for calculating $\langle N_1 \rangle$

$$\begin{aligned} \gamma_{12}^{(1)}(t) &= 2i \frac{\Omega_{24}}{\omega_{14\,12}} \sin(\omega_{14\,12}t/2) \exp(i\omega_{14\,12}t/2) \\ \alpha_{14\,34}^{(2)}(t) &= -4\Omega_{13}\Omega_{24} \left[\frac{\exp[i(\omega_{14\,34} - \omega_{14\,12})t/2] \sin[(\omega_{14\,34} - \omega_{14\,12})t/2]}{2i\omega_{14\,34}(\omega_{14\,34} - \omega_{14\,12})} - \frac{\exp(-i\omega_{14\,12}t/2) \sin(\omega_{14\,12}t/2)}{2i\omega_{14\,12}\omega_{14\,34}} \right] \end{aligned} \quad (41)$$

are obtained within time-dependent perturbation theory, considering H_0 as the initial state and switching on H_{RF} at $t = 0$. In the equation above we have defined

$$\omega_{14\,12} = U - \delta_{24} \quad (42)$$

$$\omega_{23\,12} = U - \delta_{13} \quad (43)$$

$$\omega_{14\,34} = U + \delta_{13} \quad (44)$$

$$\omega_{23\,34} = U + \delta_{24}. \quad (45)$$

Since we are interested in the time evolution of the expectation value of the particle number in state $|1\rangle$ up to second order in Ω , the only relevant second-order coefficients are $\alpha_{14\,34}^{(2)}(t)$ and $\alpha_{23\,34}^{(2)}(t)$. This is due to the fact that the states corresponding to other 2^{nd} order coefficients do not contribute to $\langle N_1 \rangle$ through the scalar product with $|12\rangle$ (0^{th} order term). The calculation of the terms $\langle\phi_I(t)|N_1|\phi_I(t)\rangle$ and $\langle\phi_{II}(t)|N_1|\phi_{II}(t)\rangle$ give the quasiparticle current (I_{ij}^s in equation (1) of the main article) when the proper normalization is taken into account.

In the term $\alpha_0\beta_0^* \langle\phi_{II}(t)|N_1|\phi_I(t)\rangle + \alpha_0^*\beta_0 \langle\phi_I(t)|N_1|\phi_{II}(t)\rangle$ we have contributions both from the scalar product between 1^{st} order terms ($\langle 14|\dots|14\rangle$)

$$\begin{aligned} \alpha_0^*\beta_0\gamma_a^{(1)*}(t)\gamma_b^{(1)}(t) + h.c. &= \\ 2 \frac{\Omega_{13}\Omega_{24}|\alpha_0\beta_0|}{\omega_{14\,12}\omega_{14\,34}} &[\cos[(\omega_{14\,34} - \omega_{14\,12})t + \varphi] + \cos(\varphi) - \cos(\omega_{14\,12}t - \varphi) - \cos(\omega_{14\,34}t + \varphi)] \end{aligned} \quad (46)$$

and from the scalar product between the 0^{th} - 2^{nd} order terms ($\langle 12|\dots|12\rangle$)

$$\begin{aligned} \alpha_0^*\beta_0 [\alpha_{14\,34}^{(2)}(t) + \alpha_{23\,34}^{(2)}(t)] + h.c. &= \\ 2\Omega_{13}\Omega_{24}|\alpha_0\beta_0| &\left[\frac{\cos[(\omega_{14\,34} - \omega_{14\,12})t + \varphi] - \cos(\varphi)}{\omega_{14\,34}(\omega_{14\,34} - \omega_{14\,12})} + \frac{\cos(\omega_{14\,12}t - \varphi) - \cos(\varphi)}{\omega_{14\,12}\omega_{14\,34}} + \right. \\ &\left. \frac{\cos[(\omega_{23\,34} - \omega_{23\,12})t + \varphi] - \cos(\varphi)}{\omega_{23\,34}(\omega_{23\,34} - \omega_{23\,12})} + \frac{\cos(\omega_{23\,12}t - \varphi) - \cos(\varphi)}{\omega_{23\,12}\omega_{23\,34}} \right]. \end{aligned} \quad (47)$$

Here φ stands for the initial phase difference between α_0 and β_0 *i.e.* $\alpha_0^*\beta_0 = |\alpha_0\beta_0|\exp(i\varphi)$. In Equations (46) and (47), the components oscillating with frequency $(\omega_{1434} - \omega_{1412})$ correspond to the contributions giving rise to the Josephson current, while components oscillating at ω_{1434} , ω_{2312} represent a contribution to the quasi-particle current, and the terms in ω_{1412} cancel out.

If we now focus on the component of $\langle N_1 \rangle$ oscillating at the Josephson frequency and replace $\omega_{ij\,kl}$ with their values in terms of interaction energies and detunings as given in equations (42) and (43), we obtain

$$\langle N_1 \rangle_J = 2\Omega_{13}\Omega_{24} \left[\frac{1}{(U + \delta_{13})(\delta_{13} + \delta_{24})} + \frac{1}{(U + \delta_{24})(\delta_{13} + \delta_{24})} + \frac{1}{(U - \delta_{24})(U + \delta_{13})} \right] \cos[(\delta_{13} + \delta_{24})t + \varphi]. \quad (48)$$

Note that we would also have a contribution to the Josephson current coming from the scalar product between the terms containing $\alpha_{1412}^{(2)}(t) + \alpha_{2312}^{(2)}(t)$ and $\alpha_{1434}^{(2)}(t) + \alpha_{2334}^{(2)}(t)$, associated with two different paths, via two different internal states, for the tunneling of Cooper pairs, but this is only a fourth order process. Such processes are present even if one of the paired states would initially be empty.

Differentiating Equation (48) with respect to time, we have

$$I_{13}^J = 2\Omega_{13}\Omega_{24}|\alpha_0\beta_0| [M_{\text{pair}} + M_{\text{single}}] \sin[(\delta_{13} + \delta_{24})t + \varphi], \quad (49)$$

where M_{pair} corresponds to the contributions to the Josephson current coming from Equation (47), *i.e.* from interference of pair tunneling and the initial population,

$$M_{\text{pair}} = \frac{1}{(U + \delta_{13})} + \frac{1}{(U + \delta_{24})} \quad (50)$$

and M_{single} to that coming from Equation (46), *i.e.* from the interference in the single-particle tunneling to the excited state |14⟩

$$M_{\text{single}} = \frac{\delta_{13} + \delta_{24}}{(U - \delta_{24})(U + \delta_{13})}. \quad (51)$$

From equation (49) we obtain the relation given in the article for the Josephson current

$$I_{13}^J = 2\Omega_{13}\Omega_{24}|\alpha_0\beta_0| \left[\frac{1}{(U + \delta_{24})} + \frac{1}{(U - \delta_{24})} \right] \sin[(\delta_{13} + \delta_{24})t + \varphi]. \quad (52)$$

Coupling dependence of the Josephson frequency

The exact numerical solution of the four-state model is very much feasible, and supplements well the perturbative analytical treatment above. In Figure 6 we show that for small couplings the perturbative treatment agrees with exact numerics as one would expect. We may then study numerically the higher order effects in the coupling strength. The dependence of the Josephson frequency on the coupling strength is a particularly interesting question. In Figure 7 we plot the value of the Josephson frequency as a function of the RF coupling. As mentioned in the article this coupling dependence of the Josephson frequency may impose a limit to the accuracy of the Josephson voltage-frequency relation.

Numerically calculated eigenstates

In general, solving analytically the eigenstates of the four state system of Figure 5 produces extremely cumbersome formulas which do not give much insight to the problem. However, to illustrate our arguments, we give here numerically calculated eigenstates in certain selected cases. Since we are now solving the eigenstates of a multiple state system where the states are coupled and have energy differences (detunings) between them, this is similar to dressed state descriptions of some quantum optics systems [1].

Zero detunings

Let us consider a system with $U/J = -5.0$. In case of zero detunings $\delta_{13}/J = \delta_{24}/J = 0$ but having finite couplings, here $\Omega_{13}/J = \Omega_{24}/J = 1$, the dressed states are the following

$$|I\rangle = -0.6672|12\rangle + 0.2341|14\rangle - 0.2341|23\rangle - 0.6672|34\rangle, \quad (53)$$

$$|II\rangle = 0.7071|12\rangle - 0.7071|34\rangle, \quad (54)$$

$$|III\rangle = 0.7071|14\rangle + 0.7071|23\rangle, \quad (55)$$

$$|IV\rangle = -0.2341|12\rangle + 0.6672|14\rangle - 0.6672|23\rangle - 0.2341|34\rangle, \quad (56)$$




Figure 6: The Josephson critical currents $|I_{13}^C|$ (dots) and $|I_{24}^C|$ (crosses) as a function of δ_{24}/J , when $\delta_{13}/J = 2.0$, given by the exact numerical calculation for the four-state system (here $U/J = 5.0$ and $\Omega/J = 0.1$). The solid lines show the results of the perturbative calculation. J is the unit of energy.




Figure 7: Position of the Josephson peak (*i.e.* the Josephson oscillation frequency) as a function of the RF coupling, with $\Omega_{13} = \Omega_{24} = \Omega$, $\delta_{13}/J = \delta_{24}/J = 1.0$ and $U/J = 10.0$.

with the eigenenergies

$$E_I = -5.7016, \quad E_{II} = -5.0, \quad E_{III} = 0.0, \quad E_{IV} = 0.7016. \quad (57)$$

In general, if the initial state of the system is one of eigenstates of the Hamiltonian (a dressed state), there will be no dynamics. Here the state $|II\rangle$ is what one could consider in our small system as an analogue of the BCS state for two superfluids: a superposition of pairs. Having that state as the initial state, no dynamics takes place; this corresponds to the absence of Josephson oscillations for zero voltage (zero detuning) in case of identical superconductors. But if the phase difference between the superconductors, or their other properties, would be such that the initial state is no longer identical to one of the eigenstates (*e.g.* $|II\rangle$), then there will be dynamics. That corresponds to the DC Josephson effect, caused by a phase difference between two superconductors instead of a voltage. This is just the standard two-mode description of Josephson physics found in many text-books (the two modes correspond to the pairs (superconductors) on both sides of the junction, here to the states $|12\rangle$ and $|34\rangle$).

Symmetric detunings

Now let us look at the effect of the detunings when they are symmetric. We take here the values $\delta_{13}/J = \delta_{24}/J = 1$. Then the eigenstates are

$$|I\rangle = 0.9581|12\rangle - 0.1737|14\rangle + 0.1737|23\rangle + 0.1470|34\rangle, \quad (58)$$

$$|II\rangle = -0.2072|12\rangle - 0.17|14\rangle + 0.17|23\rangle + 0.9483|34\rangle, \quad (59)$$

$$|III\rangle = 0.7071|14\rangle + 0.7071|23\rangle, \quad (60)$$

$$|IV\rangle = 0.1976|12\rangle + 0.664|14\rangle - 0.664|23\rangle + 0.2813|34\rangle, \quad (61)$$

with the eigenenergies

$$E_I = -6.3626, \quad E_{II} = -4.3586, \quad E_{III} = 0.0, \quad E_{IV} = 0.7212. \quad (62)$$

Now one can see that the states $|12\rangle$ and $|34\rangle$ do not appear any more symmetrically in the eigenstates. This leads to the Josephson oscillations for finite detuning (finite voltage), if we have an initial state that contains only $|12\rangle$ and $|34\rangle$, as would be the analogue for superconductors.

Note that the eigenstates contain unpaired states. These can then contribute to the dynamics. In this sense, the four state system description goes beyond the simple two-mode text-book description of the Josephson effect. Of course, these unpaired contributions are implicitly involved in more elaborate descriptions of Josephson physics, as is for instance evident from the fact that our self-consistent linear response calculation could reveal the asymmetry of the critical currents in the asymmetric detunings case.

However, in previous literature, the unpaired contributions have not been paid attention to, because, as is clear from the symmetric structure of the above eigenstates with respect to states $|14\rangle$ and $|23\rangle$, the contributions of these states will be *equal in magnitude* in the dynamics. Therefore it appears as if only pairs are tunneling and no single particles exist during the dynamics: but in fact, the single particles exist, it just happens that the expectation values of single particles in states $|14\rangle$ and $|23\rangle$ at any given time of the dynamics are the same. Therefore it seems like $|1\rangle$ and $|2\rangle$ are tunneling together. Another reason why the single particle contributions have not been notified earlier is that it is known that the standard single particle currents require detunings (voltages) above twice the gap; how could they then participate in oscillations at the below gap Josephson frequency? In this manuscript, we have shown that it is *the interference term of these single particle currents* (i.e. a beating of the standard single particle contributions) that contributes to the Josephson oscillations.

Asymmetric detunings

Finally, let us choose different detunings with $\delta_{13}/J = 1$ and $\delta_{24}/J = 2$ *i.e.* we consider the "different voltages spin up and spin down electrons" case. Then the eigenstates are

$$|I\rangle = 0.9704|12\rangle - 0.1684|14\rangle + 0.1454|23\rangle + 0.0944|34\rangle, \quad (63)$$

$$|II\rangle = -0.1582|12\rangle - 0.2291|14\rangle + 0.1774|23\rangle + 0.944|34\rangle, \quad (64)$$

$$|III\rangle = 0.0884|12\rangle + 0.9138|14\rangle + 0.3586|23\rangle + 0.1692|34\rangle, \quad (65)$$

$$|IV\rangle = -0.1599|12\rangle - 0.2902|14\rangle + 0.9049|23\rangle - 0.2672|34\rangle, \quad (66)$$

with the eigenenergies

$$E_I = -6.8234, \quad E_{II} = -3.9305, \quad E_{III} = -0.2182, \quad E_{IV} = 0.9721. \quad (67)$$

The observations of the dynamics in the previous case of symmetric detunings apply here as well, with the important difference that now the single particle contributions $|14\rangle$ and $|23\rangle$ do not appear symmetrically in the eigenstates. It is then understandable that asymmetry with respect to them appears also in the dynamics: this is the asymmetry in critical currents that we have predicted and explained. Note that by finding and explaining this asymmetry, we could provide insight also to the standard symmetric case as explained in the case of symmetric detunings: the Josephson effect consists of interference terms of pair and single particle tunneling processes, in particular, the interference of standard single particle currents contributes to the Josephson effect also in the symmetric case, although it appears as pair tunneling due to the symmetry. In the asymmetric case, the single particle interference term then clearly manifests itself, and the novel intuitive understanding of the Josephson effect proposed by us, namely interferences in pair and single particle tunnelings, can be put under a direct experimental test.

References

- [1] C. Cohen-Tannoudji, J. Dupont-Roc, and G. Grynberg, *Atom-Photon interactions* (John Wiley & Sons, 1998).
- [2] G. Baym and L. P. Kadanoff, Phys. Rev. **124**(2), 287 (1961).
- [3] G. Baym, Phys. Rev. **127**, 1391 (1962).

Spin-asymmetric Josephson effect

M.O.J. Heikkinen,¹ F. Massel,¹ J. Kajala,¹ M.J. Leskinen,¹ G.S. Paraoanu,² and P. Törmä^{1,*}

¹*Department of Applied Physics, Aalto University School of Science and Technology, P.O.Box 15100, FI-00076 Aalto, FINLAND*

²*Low Temperature Laboratory, Aalto University School of Science and Technology, P.O.Box 15100, FI-00076 Aalto, FINLAND*

We propose that with ultracold Fermi gases one can realize a spin-asymmetric Josephson effect in which the two spin components of a Cooper pair are driven asymmetrically — corresponding to driving a Josephson junction of two superconductors with different voltages $V_\uparrow$ and $V_\downarrow$ for spin up and down electrons, respectively. We predict that the spin up and down components oscillate at the same frequency but with different amplitudes. Furthermore our results reveal that the standard interpretation of the Josephson supercurrent in terms of coherent bosonic pair tunneling is insufficient. We provide an intuitive interpretation of the Josephson supercurrent as interference in Rabi oscillations of pairs and single particles, the latter causing the asymmetry.

PACS numbers: 67.85.-d, 74.50.+r, 03.75.Lm

When a coherent many-body system is partitioned into two sub-systems, the dynamics of macroscopic observables such as relative number of particles and relative phase is called the Josephson effect [1, 2]. The external Josephson effect has been realized in superconducting junctions [3, 4], superfluid ^3He [5] and ^4He [6], and in Bose-Einstein condensates (BEC) of alkali atomic gases in double-well traps [7, 8]. The internal Josephson effect has been demonstrated in ^3He [9] and is expected to occur in spin BECs [2, 10–12]. Also in the context of ultracold Fermi gases [13] the possibility of the Josephson effect has recently received theoretical interest [14–17]. In this letter we show that partitioning a system of Cooper-paired fermions so that the two components of the pair experience different potentials (this is what we mean by “spin-asymmetric” here) leads to a novel effect, namely different-amplitude but phase-synchronized number-oscillations of the components. Although the microscopic description of the Josephson effect is based on single particle tunneling, the standard interpretation of the Josephson supercurrent is given in terms of coherent tunneling of bosons or Cooper pairs [18]. Importantly, our results show that such an interpretation is insufficient. We provide a clear, intuitive explanation of the predicted spin-asymmetric Josephson effect and a new understanding of the Josephson supercurrent as a process where not only pairs but also the spin-components separately contribute via interference.

We propose that the spin-asymmetric Josephson effect can be realized in a four-component Fermi gas in which two superfluids are coupled by radio-frequency (rf) fields (Fig. 1a), as in rf spectroscopy [19]. The setup is motivated by the recent realization of three-component Fermi gases [20]. Another possible, perhaps experimentally simpler, realization is a superfluid two-component ultracold Fermi gas (Fig. 1b) in a (spin)component-dependent double-well potential. The theoretical descriptions of the systems of Fig. 1a and Fig. 1b are identical (in this letter we use the notation of the former). We also suggest

that the spin-asymmetric effect can be realized in a SIS-junction of two materials with different Zeeman splittings [21].

The setup of Fig. 1a corresponds to a many-body Hamiltonian $H = H_0 + H_{rf}$, where $H_0 = \int d\mathbf{r} \sum_i \psi_i^\dagger(\mathbf{r}) (-\frac{\nabla^2}{2m} - \mu_i) \psi_i(\mathbf{r}) + \frac{1}{2} \int d\mathbf{r} \sum_{i \neq j} U_{ij} \psi_i^\dagger(\mathbf{r}) \psi_j^\dagger(\mathbf{r}) \psi_j(\mathbf{r}) \psi_i(\mathbf{r})$. Here $\psi_i(\mathbf{r})$ and $\psi_i^\dagger(\mathbf{r})$ are the fermionic field operators for the internal state i and μ_i is the chemical potential (we assume $\mu_i \equiv \mu$), and U_{ij} give the interaction strengths in the s-wave contact potential approximation. We assume that U_{12} and U_{34} lead to pairing and set $\hbar = 1$.

In the rotating wave approximation [22] the tunneling coupling between states $|i\rangle$ and $|j\rangle$ is given by the Rabi frequency Ω_{ij} with the detuning $\delta_{ij} = \nu_{ij} - \omega_{ij}$. Here ν_{ij} is the frequency of the field and ω_{ij} is the resonance frequency of the hyperfine transition. The effect of the electromagnetic field on the system is then described by $H_{rf} = \frac{\delta_{13}}{2} \int d\mathbf{r} (\psi_1^\dagger(\mathbf{r}) \psi_1(\mathbf{r}) - \psi_3^\dagger(\mathbf{r}) \psi_3(\mathbf{r})) + \frac{\delta_{24}}{2} \int d\mathbf{r} (\psi_2^\dagger(\mathbf{r}) \psi_2(\mathbf{r}) - \psi_4^\dagger(\mathbf{r}) \psi_4(\mathbf{r})) + \Omega_{13} \int d\mathbf{r} \psi_1^\dagger(\mathbf{r}) \psi_3(\mathbf{r}) + h.c. + \Omega_{24} \int d\mathbf{r} \psi_2^\dagger(\mathbf{r}) \psi_4(\mathbf{r}) + h.c.$ In analogy to the usual Josephson junctions, the states $|1\rangle$ and $|2\rangle$ correspond to spin up and down electrons in the left side superconductor, $|3\rangle$ and $|4\rangle$ on the right. The detunings δ_{13} and δ_{24} play the role of the voltage.

The essential new feature in atomic gases is the possibility to set $\delta_{13} \neq \delta_{24}$, which in the case of the Josephson junction corresponds to spin-dependent voltages. Note that this is different from superconductor-ferromagnet-superconductor (SFS) structures [23] in which the spin-active barrier coupling plays the crucial role. Though in our case also the couplings could be different, only the spin-asymmetric potential is relevant for our predictions. Moreover, one might consider ferromagnetic superconductors [24] in a junction as a related system but those materials have most likely an exotic ground state which does not fit our description.

We now determine the transition rates (*i.e.* particle

currents) between states $|1\rangle$ and $|3\rangle$, $I_{13}(t) \equiv \langle \dot{N}_1 \rangle$, and between states $|2\rangle$ and $|4\rangle$, $I_{24}(t) \equiv \langle \dot{N}_2 \rangle$. We calculate a self-consistent linear response with respect to H_{rf} (valid when the number of transferred particles is small compared to the total particle number) for the system of Fig. 1 with the aid of the Kubo formula and the Kadanoff-Baym method [25]. We obtain the currents

$$\begin{aligned} I_{13}(t) &= I_{13}^S + I_{13}^C \sin[(\delta_{13} + \delta_{24})t + \varphi], \\ I_{24}(t) &= I_{24}^S + I_{24}^C \sin[(\delta_{13} + \delta_{24})t + \varphi]. \end{aligned} \quad (1)$$

Here I^S is the standard single particle (quasiparticle) current that occurs only for detunings δ_{ij} above the excitation gap 2Δ . The initial phase of the Josephson current is φ . The critical Josephson currents I^C become, in a spatially homogeneous case and in the BCS description,

$$I_{13}^C = 2 |\Omega_{13} \Omega_{24} \Pi_{\mathcal{F}}(\mathbf{p} = 0, \delta_{24} + i\eta^+)|, \quad (2)$$

$$I_{24}^C = 2 |\Omega_{13} \Omega_{24} \Pi_{\mathcal{F}}(\mathbf{p} = 0, \delta_{13} + i\eta^+)|, \quad (3)$$

where $\Pi_{\mathcal{F}}(\mathbf{p}, \omega) = \frac{1}{\beta V} \sum_{\mathbf{q}, \chi} \mathcal{F}_{12}(\mathbf{q}, \chi) \mathcal{F}_{34}^*(\mathbf{q} - \mathbf{p}, \chi - \omega)$. Here V is the volume and $\beta = 1/(k_B T)$ (T is temperature and k_B the Boltzmann constant). $\mathcal{F}_{12} = \mathcal{F}_{21}^*$ ($\mathcal{F}_{34} = \mathcal{F}_{43}^*$) is the anomalous mean field Matsubara Green's function for the superfluid of components $|1\rangle$ and $|2\rangle$ ($|3\rangle$ and $|4\rangle$). For details see supplementary material [26].

The striking result is that the critical current I_{13}^C in equation (2) depends only on δ_{24} , and similarly I_{24}^C in equation (3) only on δ_{13} (this is not limited to the BCS regime; it remains true whenever pairing correlations exist). By choosing different detunings δ_{13} and δ_{24} , one can observe a spin-asymmetric Josephson effect in which the currents in the two tunneling channels are different in amplitude, but oscillate at the same frequency. Moreover, the results predict a tunable DC Josephson effect: by choosing the detunings so that $\delta_{13} + \delta_{24} = 0$, the phase factor in equation (1) is constant but the critical current can still be tuned. The conclusions hold for experimentally realistic parameters and also if cross interactions (see Fig.1a) and finite temperature are included into our analysis, as shown in the supplementary material [26]. For typical parameters and taking *e.g.* $\delta_{13}/E_F = 0.4$ and $\delta_{24}/E_F = 0.5$, with E_F the Fermi energy, one obtains a considerable asymmetry of $I_{13}^C/I_{24}^C = 1.14$, see [26] for details. By performing the self-consistent calculation we have removed the ambiguity of whether our previous suggestion of the critical current asymmetry [14] was due to a simple linear response approach following the Ambegaokar-Baratoff treatment [27].

Our results (1)-(3) are in obvious contradiction with the standard interpretation of the Josephson supercurrent in terms of coherent *pair* tunneling excluding any difference in the Josephson currents of different spin-components. In what follows we provide an explanation for the spin-asymmetric Josephson effect which not only resolves this paradox but also opens




FIG. 1: Spin-asymmetric Josephson effect setup. (a) The four-component Fermi gas. Particles in internal (*e.g.* hyperfine) states $|1\rangle - |4\rangle$ coexist spatially. The components $|1\rangle$ and $|2\rangle$ as well as $|3\rangle$ and $|4\rangle$ form Cooper pairs due to the interactions U_{12} and U_{34} . The cross-interactions U_{13} , U_{14} , U_{23} , U_{24} are assumed weak, thus not leading to pairing. The tunneling is driven by applying rf fields to couple the states. The Rabi couplings Ω_{ij} correspond to the weak tunneling link between the two superconductors in a Josephson junction. The detunings $\delta_{ij} = \nu_{ij} - \omega_{ij}$ play the role of the voltage, eV . Remarkably, one can choose $\delta_{13} \neq \delta_{24}$ and create the analogue of a spin-dependent voltage in a Josephson junction. The states $|1\rangle$ and $|3\rangle$ ($|2\rangle$ and $|4\rangle$) must be states of the same atom to allow the rf coupling, but $|1\rangle$ and $|2\rangle$ can be *e.g.* ^6Li and ^{40}K . (b) Cooper pairs of a two-component Fermi gas in a spin dependent double-well potential. This setup is conceptually equivalent to the system of (a) albeit the absence of cross-interactions.

up a new point of view to the conventional Josephson effect. Let us begin by writing down the initial state of the two superfluids as a product state of two BCS states: $|\Psi\rangle = |\text{BCS}\rangle_{12} \otimes |\text{BCS}\rangle_{34} = \prod_k (u_k |0\rangle_k + v_k |12\rangle_k) \prod_{k'} (u_{k'} |0\rangle_{k'} + v_{k'} |34\rangle_{k'})$ where $|ij\rangle_k = c_{k,i}^\dagger c_{-k,j}^\dagger |0\rangle_k$. Next, we single out one Cooper pair from each superfluid. Since the rf coupling between 1-3 and 2-4 conserves momentum we focus on states with $k = k'$ (the momentum conservation can be relaxed and our conclusions still hold):

$$\begin{aligned} (u_k |0\rangle_k + v_k |12\rangle_k)(u_k |0\rangle_k + v_k |34\rangle_k) &= u_k^2 |0\rangle_k |0\rangle_k \\ &+ v_k^2 |12\rangle_k |34\rangle_k + u_k v_k |12\rangle_k |0\rangle_k + u_k v_k |0\rangle_k |34\rangle_k. \end{aligned} \quad (4)$$

The empty state $|0\rangle |0\rangle$ cannot contribute to the current and neither can $|12\rangle |34\rangle$ since it is Pauli blocked. Therefore, the Josephson physics arises from the $u_k v_k |12\rangle |0\rangle + u_k v_k |0\rangle |34\rangle$ superposition.

We then ask whether the essential features of our results can be explained by considering the dynamics of a single Cooper pair, initially in the above superposition state characteristic for the BCS state with a macroscopic phase. In addition to the paired states $|12\rangle |0\rangle \equiv |12\rangle$ and $|0\rangle |34\rangle \equiv |34\rangle$ the states $|1\rangle |4\rangle \equiv |14\rangle$ and $|2\rangle |3\rangle \equiv |23\rangle$ are required to have a closed subsystem with respect to the tunneling coupling, see Fig. 2. These broken pair states are analogous to single particle excitations within the BCS formalism.

We now solve the time evolution of this system per-

turbatively in the couplings Ω_{ij} . We consider an initial state in the general superposition form $\alpha_I |12\rangle + \beta_{II} |34\rangle$ as suggested by equation (4). The total current becomes

$$\begin{aligned} \langle \phi(t) | \dot{N}_1 | \phi(t) \rangle &= |\alpha_I|^2 \langle \phi_I(t) | \dot{N}_1 | \phi_I(t) \rangle \\ &+ |\beta_{II}|^2 \langle \phi_{II}(t) | \dot{N}_1 | \phi_{II}(t) \rangle + \alpha_I \beta_{II}^* \langle \phi_{II}(t) | \dot{N}_1 | \phi_I(t) \rangle \\ &+ \alpha_I^* \beta_{II} \langle \phi_I(t) | \dot{N}_1 | \phi_{II}(t) \rangle, \end{aligned} \quad (5)$$

where $|\phi_I(t)\rangle = \exp(-iHt) |12\rangle$ and $|\phi_{II}(t)\rangle = \exp(-iHt) |34\rangle$ are calculated to second order in Ω_{ij} . For details and for a complementary discussion in terms of exact numerical eigenstates (dressed states) see [26]. Here the first two terms contain only single particle Rabi processes, which correspond to the standard single particle (quasiparticle) currents in a Josephson junction, I_{ij}^S in equation (1). Only the last two terms in equation (5), with $\alpha_I \beta_{II}^* \equiv |\alpha_I \beta_{II}| e^{i\varphi}$, contribute to the Josephson current. Thus, the Josephson effect originates from the interference part of the Rabi oscillations in the $|12\rangle, |34\rangle, |14\rangle, |23\rangle$ state space.

Isolating the terms oscillating at the Josephson frequency in $\langle \phi(t) | \dot{N}_1 | \phi(t) \rangle$, we get the Josephson current

$$\begin{aligned} I_{13}^J &= I_{13}^C(\delta_{24}) \sin[(\delta_{13} + \delta_{24})t + \varphi], \\ I_{13}^C(\delta_{24}) &= 2\Omega_{13}\Omega_{24}|\alpha_I\beta_{II}| \left[\frac{1}{U + \delta_{24}} + \frac{1}{U - \delta_{24}} \right]. \end{aligned} \quad (6)$$

Hereby, we obtain the same qualitative result as our linear response calculation gave in equation (1): the amplitude of I_{13}^J depends only on the detuning δ_{24} . Now we can identify the source of the dependence of I_{13}^J on δ_{24} . The current I_{13}^J derives from the population of species $|1\rangle$, *i.e.* both the population of state $|12\rangle$ and $|14\rangle$. The result of equation (6) is the sum of these two contributions.

The contribution from state $|12\rangle$ is the result of tunneling between the paired states $|12\rangle$ and $|34\rangle$, via the intermediate state $|14\rangle$ (or $|23\rangle$) as shown in Fig. 3a. The term is symmetric in δ_{13} and δ_{24} and proportional to $M_{pair} = \frac{1}{U+\delta_{13}} + \frac{1}{U+\delta_{24}}$. The contribution is of second order because it originates from a zeroth order and a second order process: the population due to pair tunneling from $|34\rangle$ to $|12\rangle$ (second order) is indistinguishable from the initial (zeroth order) population in state $|12\rangle$. Pair tunneling contributions that do not originate from interference also exist but they are of fourth order in Ω .

The contribution to I_{13}^C from state $|14\rangle$ arises from interference of broken pairs (single particles) as depicted in Fig. 3b. This contribution is responsible for the asymmetry as it is proportional to $M_{single} = \frac{1}{U-\delta_{24}} - \frac{1}{U+\delta_{13}}$. Physically, this term arises because we cannot distinguish between broken pairs from the states $|12\rangle$ and $|34\rangle$. The two paths leading to the state $|14\rangle$ are not symmetric with respect to δ_{13} and δ_{24} , see Figs. 2 and 3. This gives important new insight also to the Josephson effect in general: also *single particle* interferences contribute to the Josephson current $I_{ij}^C \sin(\delta_{13} + \delta_{24})$ of equation (1).




FIG. 2: Energy level diagram of the four-state model. The basis states $|12\rangle, |34\rangle, |14\rangle, |23\rangle$ form a closed subspace under the Rabi oscillation processes caused by the rf coupling. The paired states $|12\rangle$ and $|34\rangle$ have an energy lower by U than the states of broken pairs $|14\rangle$ and $|23\rangle$.




FIG. 3: Interference processes leading to the Josephson supercurrent. The initial state is $|\phi\rangle = (|12\rangle + |34\rangle)/\sqrt{2}$. (a) The second order transition from $|34\rangle$ through $|14\rangle$ to $|12\rangle$ creates an interference with the initial (zeroth order) population in state $|12\rangle$. The resulting contribution is of the order $\Omega_{13}\Omega_{24}$. Also the same process via the state $|23\rangle$ contributes. (b) The first order transitions to the excited state $|14\rangle$ from $|12\rangle$ and $|34\rangle$ create an interference proportional to $\Omega_{13}\Omega_{24}$.

In the standard symmetric case $\delta_{13} = \delta_{24} \equiv \delta$, we have $M_{pair} = 2/(U + \delta)$ and $M_{single} = 2\delta/(U^2 - \delta^2)$. At the DC Josephson limit, $\delta \rightarrow 0$, the pair transfer M_{pair} dominates, which is intuitively appealing. Closer to the Riedel peak the excited state (single particle) interference M_{single} becomes equally important. To illustrate further, in a typical Al/AlO_x/Al junction at the voltage of 0.015 mV the single particle interference accounts for 1.7 % of the AC Josephson current. Note that our "single particle interference term" is not the cosine-term (also called "quasiparticle interference term" [28]) of the Josephson effect. The cosine-term involves real single particle transitions and exists only for voltages above 2Δ at zero temperature. In contrast, our single particle interference term is inherent in the supercurrent and corresponds to virtual transitions. Similarly, our findings are different from various combined effects of single particle currents and supercurrents in small Josephson junctions [29], where again the single particle transitions are real, not virtual. Note also that we do not consider any interactions between the Cooper pairs (analogue of charging effects) nor the effect of the environment.

We emphasise the interference nature of the Joseph-

son effect: it requires lack of which-way information on the tunneling path of the particle. Above, the superposition $u_k v_k(|12\rangle|0\rangle + |0\rangle|34\rangle)$ was due to the uncertainty of the particle number in the BCS state. It is interesting to contrast this with the number-projected BCS-state [30] or the Fock-state [31], $|\Psi\rangle_{Ncons} = (u_k v_{k'}|0\rangle_k|12\rangle_{k'} + u_{k'} v_k|12\rangle_k|0\rangle_{k'})(u_k v_{k'}|0\rangle_k|34\rangle_{k'} + u_{k'} v_k|0\rangle_{k'}|34\rangle_k)$ (for two k -states.) Here the part relevant for Josephson physics would be $u_k v_{k'} u_{k'} v_k|12\rangle_{k'}|34\rangle_k + u_k v_{k'} u_{k'} v_k|12\rangle_k|34\rangle_{k'}$. Now, the entanglement of $|34\rangle_{k'}$ with $|12\rangle_k$ allows to determine, whether a particle in state $|1\rangle$ belongs to an initial 1-2 pair or to a tunnelled 3-4 pair. The indistinguishability is lost and there is no Josephson effect. Note that the system of Fig. 1a should be cooled with the rf couplings on to realize the necessary uncertainty in particle number. One can also ask whether, in the case of the separated 1-2 and 3-4 condensates (Fock states), the measurement process itself is sufficient to generate the relative particle number uncertainty, in analogy to the famous problem of interference between two BECs [32].

Importantly, based on the picture of interfering Rabi processes, one would anticipate corrections from the coupling strength Ω_{ij} to the Josephson frequency $\delta_{13} + \delta_{24}$ given by linear response. We have observed such a correction in the numerical simulations, see supplementary material [26]. We expect this correction to vanish in the thermodynamic limit, but it could be experimentally tested in ultracold gases and in superconducting grains.

In summary, we have predicted a spin-asymmetric Josephson effect which could be observed in ultracold Fermi gases and solid state systems and is likely to be relevant for spintronics. We provide a microscopic description of the Josephson effect as interference in Rabi oscillations of pairs and single particles, where the latter cause the predicted asymmetry. In particular, our finding that the interference part of single particle currents contributes to the Josephson supercurrent, in addition to the interference in pair tunneling, is fundamental. The single particle interference contribution is present already in the standard (symmetric) case, and becomes manifest and experimentally verifiable in the asymmetric one.

We thank J.J. Kinnunen and S. van Dijken for useful discussions. This work was supported by NGSMP, FICS, EUROQUAM/FerMix and Academy of Finland (Project No. 129896, No. 135135, No. 210953, No. 213362, No. 217043, No. 217045), and conducted as a part of a EURYI scheme grant [33]. We acknowledge the use of CSC – IT Center for Science Ltd computing resources.

* Electronic address: paivi.torma@hut.fi

- [1] B.D. Josephson, *Physics Letters* **1**, 251 (1962).
- [2] A.J. Leggett, *Rev. Mod. Phys.* **73**, 307 (2001).
- [3] P.W. Anderson and J.M. Rowell, *Phys. Rev. Lett.* **10**, 230 (1963).
- [4] D.N. Langenberg *et al.*, *Phys. Rev. Lett.* **15**, 842 (1965).
- [5] S. Backhaus *et al.*, *Science* **278**, 1435 (1997).
- [6] K. Sukhatme *et al.*, *Nature* **411**, 280 (2001).
- [7] M. Albiez *et al.*, *Phys. Rev. Lett.* **95**, 010402 (2005).
- [8] S. Levy *et al.*, *Nature* **449**, 579 (2007).
- [9] R.A. Webb, R.E. Sager, and J.C. Wheatley, *Phys. Rev. Lett.* **35**, 1010 (1975); A.J. Leggett, *Rev. Mod. Phys.* **47**, 331 (1975); K. Maki and T. Tsuneto, *Prog. Theor. Phys.* **52**, 773 (1974).
- [10] D.S. Hall *et al.*, *Phys. Rev. Lett.* **81**, 1543 (1998).
- [11] P. Böhi *et al.*, *Nat. Phys.* **5**, 592 (2009).
- [12] M.-S. Chang *et al.*, *Nat. Phys.* **1**, 111 (2005).
- [13] S. Giorgini *et al.*, *Rev. Mod. Phys.* **80**, 1215 (2008); I. Bloch *et al.*, *ibid.* **80**, 885 (2008).
- [14] G.-S. Paraoanu *et al.*, *Phys. Rev. A* **66**, 041603 (2002).
- [15] R. Onofrio and C. Presilla, *Phys. Rev. A* **70**, 043608 (2004).
- [16] L. Salasnich *et al.*, *Phys. Rev. A* **77**, 043609 (2008); F. Ancilotto *et al.*, *ibid.* **79** (2009).
- [17] V.M. Galitski, *Phys. Rev. A* **72**, 013612 (2005).
- [18] P.G. de Gennes, *Superconductivity of metals and alloys* (Westview Press, Boulder, Colorado, 1999).
- [19] C.A. Regal and D.S. Jin, *Phys. Rev. Lett.* **90**, 230404 (2003); S. Gupta *et al.*, *Science* **300**, 1723 (2003); C. Chin *et al.*, *ibid.* **305**, 1128 (2004); J. Kinnunen, M. Rodriguez, and P. Törmä, *ibid.* **305**, 1131 (2004).
- [20] A.N. Wenz *et al.*, *Phys. Rev. A* **80**, 040702 (2009).
- [21] R. Meservey and P.M. Tedrow, *Phys. Rep.* **238**, 173 (1994).
- [22] C. Cohen-Tannoudji *et al.*, *Atom-photon interactions* (Wiley-Interscience, New York, 1998).
- [23] F.S. Bergeret *et al.*, *Rev. Mod. Phys.* **77**, 1321 (2005).
- [24] S.S. Saxena *et al.*, *Nature* **406**, 587 (2000).
- [25] G. Baym and L.P. Kadanoff, *Phys. Rev.* **124**, 287 (1961).
- [26] See online supplementary material.
- [27] V. Ambegaokar and A. Baratoff, *Phys. Rev. Lett.* **10**, 486 (1963).
- [28] J.R. Waldram, *Rep. Prog. Phys.* **39**, 751 (1976).
- [29] T.A. Fulton *et al.*, *Phys. Rev. Lett.* **63**, 1307 (1989); A. Maassen van den Brink, G. Schön, and L.J. Geerlings, *Phys. Rev. Lett.* **67**, 3030 (1991).
- [30] B.F. Bayman, *Nuclear Physics* **15**, 33 (1960).
- [31] A.J. Leggett, *Quantum Liquids* (Oxford University Press, Oxford, 2006).
- [32] M.R. Andrews *et al.*, *Science* **275**, 637 (1997).
- [33] See www.esf.org/euryi.